

INFORMATION TO BE GIVEN TO THE ANESTHESIOLOGIST

Polio results in widespread neural changes, not just destruction of the spinal cord anterior horn (motor nerve) cells, and these changes get worse as patients age. These anatomic changes affect many aspects of anesthesia care. No study of polio patients having anesthesia has been done. These recommendations are based on extensive review of the current literature and clinical experience with these patients. They may need to be adjusted for a particular patient.

1. Post-Polio patients are nearly always very sensitive to sedative meds, and emergence can be prolonged. This is probably due to central neuronal changes, especially in the Reticular Activating System, from the original disease.
2. Non-depolarizing muscle relaxants cause a greater degree of block for a longer period of time in Post-Polio patients. The current recommendation is to start with half the usual dose of whatever you're using, adding more as needed. This is because the poliovirus actually lived at the neuromuscular junctions during the original disease, and there are extensive anatomic changes there, even in seemingly normal muscles, which make for greater sensitivity to relaxants. Also, many patients have a significant decrease in total muscle mass. Neuromuscular monitoring intra-op helps prevent overdose of muscle relaxants. Overdose has been a frequent problem.
3. **Succinylcholine** often causes severe, generalized muscle pain postop. It's useful if this can be avoided, if possible.
4. Postop pain is often a significant issue. The anatomic changes from the original disease can affect pain pathways due to "spill-over" of the inflammatory response. Spinal cord "wind-up" of pain signals seems to occur. Proactive, multi-modal post-op pain control (local anesthesia at the incision plus PCA, etc.) helps.
5. The autonomic nervous system is often dysfunctional, again due to anatomic changes from the original disease (the inflammation and scarring in the anterior horn "spills over" to the intermediolateral column, where sympathetic nerves travel). This can cause gastro-esophageal reflux, tachyarrhythmias and, sometimes, difficulty maintaining BP when anesthetics are given.
6. Patients who use ventilators often have worsening of ventilatory function postop, and some patients who did not need ventilation have had to go onto a ventilator (including long-term use) postop. It's useful to get at least a VC preop, and full pulmonary function studies may be helpful. One group that should all have preop PFTs is those who were in iron lungs. The marker for real difficulty is thought to be a VC <1.0 liter. Such a patient needs good pulmonary preparation preop and a plan for postop ventilatory support. Another ventilation risk is obstructive sleep apnea in the postop period. Many post-polios are turning out to have significant sleep apnea due to new weakness in their upper airway muscles as they age.
7. Laryngeal and swallowing problems due to muscle weakness are being recognized more often. Many patients have at least one paralyzed cord, and several cases of bilateral cord paralysis have occurred postop, after intubation or upper extremity blocks. ENT evaluation of the upper airway in suspicious patients would be useful.
8. Positioning can be difficult due to body asymmetry. Affected limbs are osteopenic and can be easily fractured during positioning for surgery. There seems to be greater risk for peripheral nerve damage (includes brachial plexus) during long cases, probably because nerves are not normal and also because peripheral nerves may be unprotected by the usual muscle mass or tendons.

Dr. Selma H. Calmes, MD (shemd@ucla.edu) Chairman and Professor, (retired) Department of Anesthesiology, Olive View-UCLA Medical Center, Sylmar, California

This page intentionally left blank